

Product Datasheet

Prexasertib Mesylate Hydrate

Catalog No. **BP-02309B** · CAS **1234015-57-6** · Form **hydrate** · **mesylate** · Physical state **solid** · Version **1.0** · Revision **2026-09-11**


FOR RESEARCH USE ONLY. NOT FOR HUMAN OR VETERINARY USE.

Target Checkpoint Kinase (Chk); Apoptosis	Research area Cancer
---	--------------------------------

Molecular formula **C₁₉H₂₅N₇O₆S**

Molecular weight **479.5100**

STRUCTURE



DESCRIPTION

Prexasertib Mesylate Hydrate (LY2606368 Mesylate Hydrate) is a selective, ATP-competitive second-generation checkpoint kinase 1 (CHK1) inhibitor with a K_i of 0.9 nM and an IC_{50} of <1 nM. Prexasertib Mesylate Hydrate inhibits CHK2 (IC_{50} =8 nM) and RSK1 (IC_{50} =9 nM). Prexasertib Mesylate Hydrate causes double-stranded DNA breakage and replication catastrophe resulting in apoptosis. Prexasertib Mesylate Hydrate shows potent anti-tumor activity .

BIOLOGICAL ACTIVITY

In Vitro

Prexasertib Mesylate Hydrate (LY2606368 Mesylate Hydrate) inhibits MELK (IC_{50} =38 nM), SIK (IC_{50} =42 nM), BRSK2 (IC_{50} =48 nM), ARK5 (IC_{50} =64 nM). LY2606368 requires CDC25A and CDK2 to cause DNA damage.

Prexasertib Mesylate Hydrate (33, 100 nM; for 7 hours) results in DNA damage during S-phase in HeLa cells.

Prexasertib Mesylate Hydrate (8-250 nM; pre-treated for 15 minutes) inhibits CHK1 autophosphorylation (S296) and CHK2 autophosphorylation (S516) in HT-29 cells.

Prexasertib Mesylate Hydrate (4 nM; 24 hours) results in a large shift in cell-cycle populations from G1 and G2-M to S-phase with an accompanied induction of H2AX phosphorylation in U-2 OS cells.

Prexasertib Mesylate Hydrate (33 nM; for 12 hours) causes chromosomal fragmentation in HeLa cells. Prexasertib Mesylate Hydrate (100 nM; 0.5 to 9 hours) induces replication stress and depletes the pool of available RPA2 for binding to DNA.

In Vivo

Prexasertib Mesylate Hydrate (LY2606368 Mesylate Hydrate; 1-10 mg/kg; SC; twice daily for 3 days, rest 4 days; for three cycles) causes growth inhibition in tumor xenografts.

Prexasertib Mesylate Hydrate (15 mg/kg; SC) causes CHK1 inhibition in the blood and the phosphorylation of both H2AX (S139) and RPA2 (S4/S8).

REFERENCES

[1].King C, et al. LY2606368 Causes Replication Catastrophe and Antitumor Effects through CHK1-Dependent Mechanisms. *Mol Cancer Ther.* 2015 Sep;14(9):2004-13.

[2].Yin Y, et al. Chk1 inhibition potentiates the therapeutic efficacy of PARP inhibitor BMN673 in gastric cancer. *Am J Cancer Res.* 2017 Mar 1;7(3):473-483.

Tel: +86 20 31130026 · Fax: +86 20 36822386 · tech@biochemprobe.com · www.biochemprobe.com

BiochemProbe is the catalog brand of Guangzhou Yuyuan Biotech Co., Ltd. · For research use only.

Lot CoA and SDS are separate documents.